

Original Article

Decreasing Discomfort During Shock-Wave Lithotripsy Using Transcutaneous Electrical Nerve Stimulation

Fateme Guitynavard¹, Mohaddeseh Azadvari², Maryam Gholamnejad¹, Mehdi Naghdipoor Farhadabad¹, Leila Zareian Baghdadabad¹, Alireza Gorji Daroonkolae¹, Rouhollah Afsari¹, Elham Madreseh¹, Ramin Rahimnia^{*3}

¹*Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran*

²*Department of Physical Medicine and Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran*

³*Department of Medical Nanotechnology, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran*

HIGHLIGHTS

- TENS can be suggested as effective and safe pain relief during ESWL.
- One of the problematic issues during Extracorporeal Shock Wave Lithotripsy (ESWL) is pain because anesthesia can have some side effects.
- we evaluate the Transcutaneous Electric Nerve Stimulation (TENS) as a non-invasive analgesic strategy during ESWL.

ARTICLE INFO

Receive Date: 07 May 2022

Accept Date: 20 June 2022

Available online: 21 June 2022

DOI: 10.22034/TRU.2022.341228.1111

*Corresponding Author:

Ramin Rahimnia

Email: rrahimnia@tums.ac.ir

Address: Department of Medical Nanotechnology, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran.

Introduction

Urolithiasis is one of the most commonly encountered urological problems, with a lifetime prevalence of 1% to 15%, varying according to age, gender, race, and geographic location. Since the 1980s, Extracorporeal

ABSTRACT

Introduction

Shock wave lithotripsy (SWL) is a less-invasive procedure for treating a subgroup of renal stones. Since it may cause significant pain and anxiety during the procedure, several strategies have been proposed to reduce the discomfort during the SWL procedure. This study aimed to evaluate transcutaneous electrical nerve stimulation (TENS) as a non-invasive analgesic strategy during SWL.

Methods

A total of seventy-nine patients who underwent the SWL were included. The participants were randomly divided into two groups: the case group using conventional TENS (41 patients) and the control group without TENS. The visual analog scale (VAS) was recorded at the end of the SWL session. The analgesic (30mg ketorolac) was given in both groups due to the patient's request.

Results

Both study groups were similar in age, sex, BMI, history of SWL, hypertension, stone location, and stone size. Our data indicated less VAS in the case group than in the control group throughout the procedure, although it was not statistically significant (P-value=0.087). Conclusion: TENS can be offered as a non-invasive and safe pain relief strategy during the SWL.

Conclusions

TENS can be offered as a non-invasive and safe pain relief strategy during the SWL.

Keywords: Extracorporeal Shock Wave Lithotripsy; Transcutaneous Electrical Nerve Stimulation, Urolithiasis; Pain

Shock Wave Lithotripsy (SWL) has become one of the first-line options for treating some renal and ureteral calculi. Even though the method is considered a less invasive technique, pain caused by shockwaves is still a significant concern (1-3). The pain can be the consequence of two

factors: First, the increased pressure within the kidney, and second, the SWL-related trauma of the skin and muscles. Furthermore, the effectiveness of this procedure is strongly linked to SWL-related pain and anxiety (4, 5). Pain during SWL may lead to a higher rate of kidney hematomas due to a blood pressure increase (6). Several medications and methods, such as local, intravenous (IV), or oral non-steroidal anti-inflammatory drugs (NSAIDs) and opioid agents, are considered for pain management during SWL (7). The SWL has been widely used for treating urinary tract stones and is routinely administered as an outpatient procedure, although some patients do not tolerate it without analgesia (8). Transcutaneous electrical nerve stimulation (TENS) is a potential noninvasive pain relief strategy. In this method, low voltage electrical impulses transition through electrodes attached to the skin over the painful area. It is usually used to relieve a variety of painful conditions. A TENS unit contains an electrical signal generator, a battery, and a set of electrodes (9). Its mechanism is stranded by the 'gate control theory of pain transmission. The analgesic effects arise from the electrical stimulation of non-noxious afferent nerve fibers in the skin, inhibiting the transmission of nociceptive responses through the dorsal horn of the spinal cord. This effectively 'closes the gate', thereby decreasing the transmission of pain signals from the source of acute pain in the periphery to the central nervous system (10).

This article aimed to investigate the effectiveness of TENS, compared with 'sham' (i.e., placebo) to reduce pain in SWL.

Methods

The study was approved by the ethical committee institute in Iran (IR.TUMS.VCR.REC.1398.1041), and written informed consent was signed by patients. This prospective, randomized, comparative clinical study was done in the SWL unit in the Sina hospital and included 79 patients scheduled for SWL. All patients underwent a baseline evaluation of the renal function, urine analysis, urine culture, serum creatinine, and an abdominopelvic computed tomography (CT). The patient characteristics, including age, sex, body mass index (BMI), history of hypertension, SWL, and the size and location of the stone, were also recorded.

The inclusion criteria were all patients with 6–20 mm renal stones. The exclusion criteria were coagulation disorder, uncontrolled hypertension, uncontrolled arrhythmia, abdominal aortic aneurysm >4cm, pregnancy, demand pacemaker, dermatological lesions at the site of SWL (e.g., eczema or dermatitis), drug or alcohol addiction, stone in a solitary kidney, congenital urogenital anomaly, American Society of Anesthesiologists (ASA) physical status classification > II, age <18 and >80 years, multiple renal stones, history of infectious or struvite

stone, stents such as double J (DJ) in urinary tract system, patients with immunodeficiency, flank pain (VAS >0). Seventy-nine patients were allocated to case (TENS) and control groups based on block randomization with blocks of size 6. All patients received 500–1000 ml of Ringer acetate solution over 30 min during the procedure. No premedication was given.

Case Group: Forty Patients in this group were informed about the TENS application before the procedure. One channel was used, in which two stimulator electrodes were applied paravertebrally just above and below the lithotripter shock tube. A stimulation frequency of 80 Hz and pulse width of 250 μ s was selected. The current intensity was started at 10 mA and increased gradually until the patient perceived a prickling sensation with no pain. At that moment, shock waves were started pulse width (duration), 250-microsecond pulses for 5 minutes in a 10-seconds-on 20-seconds-off duty cycle.

Control Group: Thirty-eight patients with sham TENS.

In both groups number of shock waves was 2500–3000 shocks, with a frequency of 60–90/min, and voltage intensity was started at 1 kV and then increased gradually until a maximum of 16 kV. Pain scores were assessed with the 10-score linear Visual Analogue pain Scale (VAS). Parenteral ketorolac (30 mg) was given as the patient preference. At the end of each session, a questionnaire concerning the pain experience was completed by all patients using a visual analog scale (VAS; 0 no pain, 10 unbearable pain).

Statistical analysis

Baseline demographic and clinical characteristics are described as median (IQR) or frequency (percentage %) and compared between case and control groups. The Chi-square, Mann-Whitney U Test, and ordinal logistic regression were performed using R.4.0.2 and IBM SPSS (version 23; SPSS Inc., Chicago, IL, USA). Two-tailed (P values < 0.05) were considered significant.

Results

Out of 79 patients, 41 were in the case group, including 68% men (28) and 32% women (13) with a median age of 46 (IQR:33–54) years. In the control group, 38 patients were enrolled, consisting of 73% men (29) and 27% women (11) with a median age of 41 (IQR:35–49) years. The participants' history of SWL and hypertension were 75% and 21%, respectively. The mean \pm SE of the VAS score was 4.45 \pm 0.18. The size and location of the stones were similar in both groups (P-value >0.05).

There were no significant differences between both study groups in terms of age, sex, BMI, history of SWL, hypertension, stone location, and stone size (Table 1). Figure 1 showed that the mean VAS score in the control group (4.8 \pm 0.27) was higher than the case

Table 1. Comparing baseline and clinical characteristics of study subjects between TENS and control groups

Variables	Total (n=79)	Control Group (n=38)	TENS Group (n=41)	P-value	
Sex (male), n (%)	55(69.6%)	27(71.1 %)	28(68.3 %)	0.678	
Age (years), median (IQR)	42(33-53)	41(33-51)	46 (33-54)	0.743	
BMI (Kg/m2), median (IQR)	28(25-31)	28(25-31)	27(24-31)	0.582	
History of stone crushing (yes), n (%)	59(74.7%)	27(71.1 %)	32(78.0 %)	0.698	
Location (right), n (%)	39(49.4%)	20(52.6 %)	19(46.3 %)	0.651	
Hypertension (yes), n (%)	16(20.3%)	8(22.1 %)	8(19.5 %)	0.739	
Stone size (mm), n (%)	6-10	31(39.2%)	16(42.1 %)	15(36.6 %)	0.599
	11-15	32(40.5%)	13(34.2 %)	19(46.3 %)	
	16-30	13(16.5%)	7 (18.4 %)	6(14.6 %)	

IQR: Inter quartile range; Chi-squared tests were used for association between categorical variable; and Man-Whitney U Tests for comparing continues variables between groups, Because of missing value, sum of percentage may not be equal to 100%.

Table 2. Pain intensity (VAS) in the end of study according to baseline characteristics

Variables	Pain intensity (VAS)				OR (95%CI)	P-value
	Mild (n=15)	Moderate (n=35)	Severe (n=25)	Very Severe (n=4)		
TENS	10(24%)	19(46%)	11(27%)	1(2%)	0.48(0.21,1.10)	0.087
Control (reference)	5(13%)	16(42%)	14(37%)	3(8%)		

VAS: Visual analog scale; values in each cell are frequency (Row percentage); OR: odds ratio; OR and P-value were calculated based on Ordinal Regression Model.

group (4.1±0.25) but this difference was not statistically significant (P-value>0.088).

In Table 2, pain intensity is classified as mild pain (VAS: [0-2]), moderate pain (VAS: [3-5]), severe pain (VAS:[5-7]), and very severe pain (VAS: [8-10]). The frequency distribution of each category showed that many patients in the case group had mild, moderate, and severe pain, but many patients in the control group had moderate and severe pain. In the case group, 10 patients (24%) experienced mild pain, 19 patients (46%) experienced moderate pain, 11 patients (27%) experienced severe pain, and only one patient experienced very severe pain. However, in the control group, 5 patients (13%) experienced mild pain, 16 patients (42%) experienced moderate pain, 14 patients (37%) experienced severe pain, and 3 patients (8%) experienced very severe pain. Finally, we used ordinal logistic regression to detect any differences in pain severity between case and control groups. It was observed that patients in the case group had about 50% less pain compared to the control group, but it was not statistically significant.

Discussion

Treatment of urolithiasis has been revolutionized with extracorporeal shock wave lithotripsy (SWL) due to its simplicity, non-invasive nature, efficacy, and

minimal morbidity (11-13). Pain experienced during SWL is considered multifactorial, including the type of lithotripter used, frequency, voltage, age, and sex of the patient (14). There is no consensus on the standard analgesic regimen for controlling pain during SWL. Conventionally, NSAIDs, opioids, and local anesthetics are used frequently (15). Transcutaneous electrical nerve stimulation (TENS) is a non-pharmacological, non-invasive, and inexpensive analgesic technique that has been practiced for pain control in many clinical settings (16). Previous studies have also reported that TENS can be used efficiently during SWL (17).

This article aimed to investigate the effectiveness of TENS, compared with 'sham' (i.e., placebo) to reduce pain in SWL.

The results of our study have similarities and differences with studies that have used TENS to manage acute pain. Karamaz and colleagues defined which TENS modality (conventional versus acupuncture-like) is more effective as an additional analgesic regimen during extracorporeal shock-wave lithotripsy (SWL). They proved that conventional TENS was more effective than acupuncture-like TENS in reducing alfentanil consumption and alfentanil-related side effects (18).

In the study, Randa Ali Shoukry and colleagues stated that TENS is an effective and safe practice in controlling




Figure 1. Comparing mean VAS score between TENS and Control Group

pain during SWL, as it provides good analgesia and tolerability. It leads to a decrease in analgesic consumption and its side effects, with greater patient satisfaction. It decreases the time to discharge compared to fentanyl, which is ideal for outpatient procedures (9). In the study, O Reichelt and colleagues stated that TENS is a non-invasive, cost-effective method to achieve side-effect-free analgesia in SWL using third-generation lithotripters (19). Finally, due to the study's limitations and the small sample size, it is recommended that a similar study with a larger sample size be conducted more comprehensively. It is also recommended to examine the subgroups related to the demographic characteristics of patients to determine the differences in demographic variables on the effectiveness of TENS.

Conclusions

TENS can be considered an effective and safe pain relief during SWL although not established in this study.

Authors' contributions

All authors contributed equally.

Acknowledgments

Special thanks to the Urology Research Center at Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran.

Conflict of interest

All authors declare that there is no potential competing or conflict of interest.

Funding

There was no funding.

Ethics statement

This study design were approved by the Ethical Committee

of Tehran University of Medical Sciences (IR.TUMS.VCR.REC.1398.1041).

Data availability

Data will be provided on request.

Abbreviations

ASA	American Society of Anesthesiologists
BMI	Body mass index
CT	Computed tomography
NSAID	Non-steroidal anti-inflammatory drugs
SWL	Shock wave lithotripsy
TENS	Transcutaneous electrical nerve stimulation
VAS	Visual analog scale

References

1. Laksita TB, Soebadi MA, Wirjopranoto S, Hidayatullah F, Klop-ing YP, Rizaldi F. Local anesthetics versus systemic analgesics for reducing pain during Extracorporeal Shock Wave Lithotripsy (ESWL): A systematic review and meta-analysis. *Turk J Urol.* 2021;47(4):270-8.
2. Ucer O, Ceylan Y, Ekren F, Ozan E, Muezzinoglu T. Effect of anxiety and pain on success of shockwave lithotripsy (SWL) for treatment of proximal ureteral and renal pelvic stones. *Urolithiasis.* 2016;44(6):559-64.
3. Aghamir SMK, Elmimehr R, Modaresi SS, Salavati A. Comparing bleeding complications of double and single access totally tubeless PCNL: is it safe to obtain more accesses? *Urologia internationalis.* 2016;96(1):73-6.
4. Wang Z, Chen G, Wang J, Wei W. Eutectic mixture of local anaesthetics for pain reduction during extracorporeal shockwave lithotripsy: A systematic review and meta-analysis. *Plos one.* 2020;15(10):e0237783.
5. Mohseni M, Khazaeli MH, Aghamir SMK, BINIAZ A. Changes in intrarenal resistive index following electromagnetic extracorporeal shock wave lithotripsy. 2007.
6. Tauzin-Fin P, Delort-Laval S, Krol-Houdek M-C, Maurette P, Bannwarth B. Effect of balanced analgesia with buprenorphine on pain response and general anaesthesia requirement during lithotripsy procedures. *European journal of anaesthesiology.* 1998;15(2):147-52.
7. Fouladi A, Soleimani A. Comparison of different analgesic techniques for pain relief during extracorporeal shock wave lithotripsy: a double-blind, randomized clinical trial. *Acta Informatica Medica.* 2017;25(2):94.
8. Yayik AM, Ahiskalioglu A, Alici HA, Celik EC, Cesur S, Ahiskalioglu EO, et al. Less painful ESWL with ultrasound-guided quadratus lumborum block: a prospective randomized controlled study. *Scandinavian journal of urology.* 2019;53(6):411-6.
9. Shoukry RA, Al-Ansary AM. Transcutaneous electric nerve stimulation (TENS) for pain relief during extracorporeal shock-wave lithotripsy (ESWL). *Egyptian Journal of Anaesthesia.* 2019;35(1):71-6.
10. Simpson PM, Fouche PF, Thomas RE, Bendall JC. Transcutaneous electrical nerve stimulation for relieving acute pain in the prehospital setting: a systematic review and meta-analysis of randomized-controlled trials. *European Journal of Emergency Medicine.* 2014;21(1):10-7.
11. Chaussy C, Brendel W, Schmiedt E. Extracorporeally induced destruction of kidney stones by shock waves. *The Lancet.* 1980;316(8207):1265-8.
12. Aghamir SMK, Khorrami MH, Saatchi M, Zia H, Seyedesmaeli SN. A Comparison of Minimally Invasive Surgery Ureterolithotomy and Transurethral Lithotripsy in Combination with Retrograde Intrarenal Surgery: A Randomized Clinical Trial. *Translational Research in Urology.* 2020;2(4):127-31.
13. Fakhri Yasser A, Hamidi M, Aghamir SMK. Comparison of Stone-Free Rates after Ureteroscopic Pneumatic Lithotripsy in Impacted vs. Non-Impacted upper Ureteral Stones. *Translational Research in Urology.* 2019;1(2):79-83.
14. Salinas A, Lorenzo-Romero J, Segura M, Calero M, Hernández-Millán I, Martínez-Martín M, et al. Factors determining analgesic and sedative drug requirements during extracorporeal shock wave lithotripsy. *Urologia internationalis.* 1999;63(2):92-101.
15. Bach C, Zaman F, Kachrilas S, Kumar P, Buchholz N, Masood J. Drugs for pain management in shock wave lithotripsy. *Pain research and treatment.* 2011;2011.
16. Tang Z-Y, Wang H-Q, Xia X-L, Tang Y, Peng W-W, Hu L. Mechanisms and applications of transcutaneous electrical nerve stimulation in analgesia. *Sheng li xue bao:[Acta physiologica Sinica].* 2017;69(3):325-34.
17. Torres W, Fraser N, Baumgartner B, Nelson R, Evans G, Jones V, et al. The use of transcutaneous electrical nerve stimulation during the biliary lithotripsy procedure. *The Journal of stone disease.* 1992;4(1):41-5.
18. Karamaz A, Kaya S, Karaman H, Turhanoglu S. Effect of the frequency of transcutaneous electrical nerve stimulation on analgesia during extracorporeal shock wave lithotripsy. *Urological research.* 2004;32(6):411-5.
19. Reichelt O, Zermann D-H, Wunderlich H, Janitzky V, Schubert J. Effective analgesia for extracorporeal shock wave lithotripsy: transcutaneous electrical nerve stimulation. *Urology.* 1999;54(3):433-6.

Author (s) biosketches

Guitynavard F, Assistant Professor, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran.

Email: f_guitynavard@ymail.com

Azadvari M, MD, Department of Physical Medicine and Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran.

Email: drazadvari@yahoo.com

Gholamnejad M, MD, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran.

Email: mary_medical86@yahoo.com

Naghdipoor Farhadabad M, MD, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran.

Email: naghdipoormehdi@gmail.com

Zareian Baghdadabad L, MSc, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran.

Email: l-zareian@farabi.tums.ac.ir

Gorji Daroonkolaee A, MD, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran.

Email: Gorjialireza9@gmail.com

Afsari R, MD, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran.

Email: dr.afsari@gmail.com

Madreseh E, PhD, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran.

Email: emadreseh@yahoo.com

Rahimnia R, Assistant Professor, Department of Medical Nanotechnology, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran.

Email: rrahimnia@tums.ac.ir

How to cite this article

Guitynavard F, Azadvari M, Gholamnejad M, Naghdipoor Farhadabad M, Zareian Baghdadabad L, Gorji Daroonkolaee A, Afsari R, Madreseh E, Rahimnia R. Decreasing Discomfort during Shock-Wave Lithotripsy Using Transcutaneous Electrical Nerve Stimulation. *Translational Research in Urology*. 2022 Jun;4(2):71-76.

DOI:10.22034/TRU.2022.341228.1111

URL: https://www.transresurology.com/article_152050.html

